



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# A NOTE ON THE PRECEDING PAPER \*

BY

D. R. CURTISS

The results of Dr. Walsh's paper are so interesting and capable of so many applications that it may be worth while to indicate a briefer proof of his fundamental theorem which connects with earlier work in this field.

First let us note that Walsh's Theorem II is a form of Laguerre's Theorem given below. In fact, if we write

$$f(z) = (z - \alpha_1)(z - \alpha_2) \cdots (z - \alpha_k),$$

Walsh's relation

$$\frac{1}{z - \alpha_1} + \frac{1}{z - \alpha_2} + \cdots + \frac{1}{z - \alpha_k} = \frac{k}{z - \alpha}$$

becomes

$$\frac{f'(z)}{f(z)} = \frac{k}{z - \alpha},$$

or

$$(1) \quad \alpha = z - k \frac{f(z)}{f'(z)}.$$

Walsh's Theorem II states that if the roots of  $f(z)$  are in the circular region  $C$ , and if  $z$  is exterior to  $C$ , then  $\alpha$  lies in  $C$ . This is precisely the result obtained by Laguerre as given on page 59 of volume I of his collected works, if expressed in non-homogeneous coördinates as on page 57.

We shall use another form of Laguerre's Theorem (loc. cit., p. 57) to prove the underlying proposition on which Walsh bases his proof of his Theorem I, as follows:

*Every circle through any point  $z$  and its "derived point"  $\alpha$  as defined by (1) either passes through all the roots of  $f(z)$ , or else has at least one root in the region interior to it, and at least one root in the exterior region.*

I have recently called attention<sup>†</sup> to the fact that this is a corollary of a theorem of Bôcher's on jacobians which has served as a starting point in Walsh's earlier papers.

With a slight change in Walsh's notation, whereby we substitute  $z$ 's for  $\alpha$ 's, the proposition from which Theorem I is deduced may be stated thus:

---

\* Presented to the Society April 14, 1922, under the title *On the zeros of successive polars of a binary form*.

† Science, vol. 55 (1922), p. 193.

Let  $z_1, z_2, \dots, z_k$  lie within a circular region  $C$ , and let  $\sum_r^k$  denote the sum of all products of the  $z$ 's taken  $r$  at a time. Then the equation

$$(2) \quad a_0 z^k + k a_1 z^{k-1} + \frac{k(k-1)}{2!} a_2 z^{k-2} + \dots + k a_{k-1} z + a_k = 0,$$

where

$$-a_k = a_0 \sum_k^k + a_1 \sum_{k-1}^k + a_2 \sum_{k-2}^k + \dots + a_{k-1} \sum_1^k$$

has at least one root in  $C$  for each set of values  $a_0, a_1, \dots, a_{k-1}$ .

This theorem is trivial if all the  $a$ 's are zero. In any other case we proceed by induction, assuming our proposition true for  $k = r - 1$  and proving its validity for  $k = r$ . It is obviously true for  $k = 1$ .

Let us place  $k = r$  in (2) and in (1). The resulting left-hand member of (2) we will use as  $f(z)$  in (1). We write

$$f(z) = a_0 z^r + r a_1 z^{r-1} + \dots + a_r,$$

$$zf'(z) - rf(z) = -r[a_1 z^{r-1} + (r-1)a_2 z^{r-2} + \dots + a_r],$$

so that (1) becomes

$$(3) \quad \alpha = -\frac{a_1 z^{r-1} + (r-1)a_2 z^{r-2} + \dots + a_r}{a_0 z^{r-1} + (r-1)a_1 z^{r-2} + \dots + a_{r-1}}.$$

Now consider the equation obtained by replacing  $\alpha$  in (3) by  $z_r$ ; cleared of fractions and arranged according to powers of  $z$  this is

$$(4) \quad (a_0 z_r + a_1) z^{r-1} + (r-1)(a_1 z_r + a_2) z^{r-2} + \dots + (a_{r-1} z_r + a_r) = 0.$$

But if we compare this with (2), and note that

$$\begin{aligned} -(a_{r-1} z_r + a_r) &= -a_{r-1} z_r + a_0 \sum_r^r + a_1 \sum_{r-1}^r \\ &\quad + a_2 \sum_{r-2}^r + \dots + a_{r-1} \sum_1^r \\ &= -a_{r-1} z_r + a_0 z_r \sum_{r-1}^{r-1} + a_1 (z_r \sum_{r-2}^{r-1} + \sum_{r-1}^{r-1}) \\ &\quad + a_2 (z_r \sum_{r-3}^{r-1} + \sum_{r-2}^{r-1}) + \dots + a_{r-1} (z_r + \sum_1^{r-1}) \\ &= (a_0 z_r + a_1) \sum_{r-1}^{r-1} + (a_1 z_r + a_2) \sum_{r-2}^{r-1} \\ &\quad + \dots + (a_{r-2} z_r + a_{r-1}) \sum_1^{r-1}, \end{aligned}$$

we see that (4) is a case of (2) for  $k = r - 1$ , and hence (4) has a root  $Z_r$  in  $C$ .

We now give  $z$  the value  $Z_r$  in (3) and obtain the result that the "derived point"  $\alpha$  is  $z_r$ . By Laguerre's Theorem every circle through  $Z_r$  and  $z_r$  either has on it a root of  $f(z)$  or has a root in its interior and one exterior to it; but  $Z_r$  and  $z_r$  are in  $C$ , so that we can draw through them a circle interior to  $C$ . Thus  $f(z)$  has a root in  $C$ , and on completing our induction we see that (2) has a root in  $C$ .

The theorem we have just proved can not, however, be regarded as a new one. It is equivalent to the theorem of Grace:\*

*If a form  $f(z)$  is apolar to a given form  $\phi(z)$ , then it has a zero within any circle enclosing all the roots of  $\phi(z)$ .*

If equation (2) and the equation

$$(5) \quad (z - z_1)(z - z_2) \cdots (z - z_k) = 0$$

are made homogeneous, the condition that (2) and (5) be apolar is precisely the relation just following (2). Since this holds, Grace's theorem requires that a root of (2) lie within any circle enclosing all the roots of (5), which are  $z_1, z_2, \dots, z_k$ .

Grace's theorem is a consequence of Laguerre's when the latter is put in homogeneous form. If in (1) we replace  $\alpha$  by  $\xi/\eta$ ,  $z$  by  $x/y$ , and  $f(z)$  by  $y^k f(x, y)$ , (1) takes the form

$$(6) \quad \xi f'_x + \eta f'_y = 0,$$

which is the equation of the first polar of  $(\xi, \eta)$  with respect to  $f(x, y)$ . One form of Laguerre's theorem (loc. cit., p. 59) may, then, be thus stated:

*If the point  $\alpha = \xi/\eta$  is without a circle  $C$  containing all the roots of  $f$ , all the roots of the first polar of  $\alpha$  with respect to  $f$  are within  $C$ .*

Consider a set of points  $z_1, z_2, \dots, z_k$ , and let  $f_1$  be the first polar of  $z_1$  with respect to  $f$ , let  $f_2$  be the first polar of  $z_2$  with respect to  $f_1$ , etc. If all these  $z$ 's are outside  $C$ , the roots of all the polynomials  $f_1, f_2, \dots, f_{k-1}$  are within  $C$ . If (2) and (5) are apolar,  $z_k$  is the root of  $f_{k-1}$ , and Grace's theorem, in which  $f$  and  $\phi$  may be interchanged, follows directly.

The "mixed polars"  $f_1, f_2, \dots$  are perhaps worth writing down in another form. Thus if we use the notation, for  $r < k$ ,

$$\psi(z) = (z - z_1)(z - z_2) \cdots (z - z_r) = c_0 z^r + c_1 z^{r-1} + \cdots + c_r,$$

we have

$$(-1)^r f_r(x, y) = c_r \frac{\partial^r f}{\partial x^r} - c_{r-1} \frac{\partial^r f}{\partial x^{r-1} \partial y} + \cdots + (-1)^r c_0 \frac{\partial^r f}{\partial y^r}.$$

In non-homogeneous notation the equation  $f_r = 0$  becomes

---

\* *The zeros of a polynomial*, Proceedings of the Cambridge Philosophical Society, vol. 11 (1900-02), p. 352. See also Szegő's paper, *Bemerkungen zu einem Satz von J. H. Grace über die Wurzeln algebraischer Gleichungen*, Mathematische Zeitschrift, vol. 13 (1922), p. 28, in which the "Faltungssatz" is, in a sense, the complement of the theorem proved above.

$$\begin{aligned}
 (7) \quad & \psi \frac{d^r f}{dz^r} - (k - r + 1) \frac{d\psi}{dz} \frac{d^{r-1} f}{dz^{r-1}} + \frac{(k - r + 1)(k - r + 2)}{2} \frac{d^2 \psi}{dz^2} \frac{d^{r-2} f}{dz^{r-2}} \\
 & + \cdots + (-1)^r \frac{(k - r + 1)(k - r + 2) \cdots k}{r!} \frac{d^r \psi}{dz^r} f = 0.
 \end{aligned}$$

*If every root of  $f$  is within a circle  $C$ , and every root of  $\psi$  is without  $C$ , then every root of (7) is within  $C$ .*

NORTHWESTERN UNIVERSITY,  
EVANSTON, ILL.